

# PROBABILITY-ONE HOMOTOPY METHODS FOR NONSMOOTH NONLINEAR SYSTEMS OF EQUATIONS

Layne T. Watson  
Departments of Computer Science and Mathematics  
Virginia Polytechnic Institute and State University  
Blacksburg, VA 24061-0106 USA

<http://www.cs.vt.edu/~ltw/>



# OUTLINE

1. Background—a homotopy methods tutorial.
2. Software—HOMPACK90, POLSYS\_PLP.
3. Application to circuit design.
4. Nonsmooth functions—definitions.
5. Probability-one homotopy theory for nonsmooth functions.
6. Application—mixed complementarity problem.

# BACKGROUND

Problem: given  $C^2 F : E^n \rightarrow E^n$ , solve  $F(x) = 0$ .

*Definition.* Let  $U \subset E^m$  and  $V \subset E^n$  be open, and  $\rho : U \times [0, 1) \times V \rightarrow E^n$  be  $C^2$ .  $\rho$  is said to be *transversal to zero* if  $D\rho$  has full rank on  $\rho^{-1}(0)$ .

*Parametrized Sard's Theorem.* Let  $\rho : U \times [0, 1) \times V \rightarrow E^n$  be  $C^2$  and define

$$\rho_a(\lambda, x) = \rho(a, \lambda, x).$$

If  $\rho$  is transversal to zero, then for almost all  $a \in U$ ,  $\rho_a$  is also transversal to zero.


Typical choices for  $\rho_a$ :

$$\rho_a(\lambda, x) = \lambda F(x) + (1 - \lambda)(x - a),$$

$$\rho_a(\lambda, x) = \lambda F(x) + (1 - \lambda)G(x; a),$$

$$\rho_a(\lambda, x) = F(x) - (1 - \lambda)F(a).$$

# BACKGROUND



Typical  $\rho_a^{-1}(0)$  for  $\rho_a(\lambda, x)$  transversal to zero.

## GLOBALY CONVERGENT PROBABILITY-ONE HOMOTOPY ALGORITHM

Given open  $U \subset E^m$ , open  $V \subset E^n$ , and  $C^2$   $F : V \rightarrow E^n$ , choose  $\rho : U \times [0, 1) \times V \rightarrow E^n$  such that

(1)  $\rho(a, \lambda, x)$  is  $C^2$  and transversal to zero, and

for each  $a \in U$ ,

(2)  $\rho_a(0, x) = 0$  has a unique solution  $x^a$  at which  $\text{rank } D_x \rho_a(0, x^a) = n$ ,

(3)  $\rho_a(1, x) = F(x)$ ,

(4)  $\rho_a^{-1}(0)$  is bounded.

Then for almost all  $a \in U$  there exists a zero curve  $\gamma$  of  $\rho_a$ , along which  $D\rho_a$  has full rank, emanating from  $(0, x^a)$  and reaching a point  $(1, \bar{x})$  where  $F(\bar{x}) = 0$ .  $\gamma$  has finite arc length if  $DF(\bar{x})$  is invertible.  $\gamma$  does not intersect itself, and is disjoint from all other zeros of  $\rho_a$ .

# SOFTWARE — HOMPACK90

- Fortran 90 modules.
- Specialized code for  $x = f(x)$ ,  $F(x) = 0$ , and  $\rho(a, \lambda, x) = 0$ .
- Three different curve tracking algorithms (ODE based, normal flow, augmented Jacobian matrix).
- Special algorithms for small dense and large sparse Jacobian matrices.
- Easy to use code for polynomial systems.

# SOFTWARE — POLSYS\_PLP

- For polynomial systems with complex coefficients.
- Implements state-of-the-art theory for exploiting structure.
- Elegant interface for defining system and its structure.
- Fortran 90 modules.

# CIRCUIT DESIGN APPLICATION

Variable stimulus homotopy:

$$\rho_a(\lambda, x) = \lambda F(x, \lambda) + (1 - \lambda)G(x, a).$$

$F(x, \lambda)$  turns on nonlinear circuit components as  $\lambda$  increases from 0 to 1.  $G(x, a)$  models a simple linear circuit.  $\rho_a(\lambda, x)$  models **some** no gain circuit for all  $0 \leq \lambda \leq 1$ .

Reference: R. C. Melville, Lj. Trajković, S.-C. Fang, and L. T. Watson, “Artificial parameter homotopy methods for the DC operating point problem”, *IEEE Trans. Computer-Aided Design*, 12 (1993) 861–877.

# NONSMOOTH FUNCTIONS

Let  $F : E^n \rightarrow E^n$  be locally Lipschitzian (hence differentiable on a dense set  $D_F$ ). The  $B$ -subdifferential is

$$\partial_B F(x) := \left\{ V \mid \exists \{x^k\} \rightarrow x, x^k \in D_F, \text{ with } V = \lim_{k \rightarrow \infty} \nabla F(x_k) \right\}.$$

The Clarke subdifferential  $\partial F(x)$  is the convex hull of  $\partial_B F(x)$ .  $F$  is said to be *semismooth* at  $x$  if it is directionally differentiable at  $x$  and for any  $V \in \partial F(x+h)$ ,  $h \rightarrow 0$ ,

$$Vh - F'(x; h) = o(\|h\|).$$

$F$  is said to be *strongly semismooth* if additionally,

$$Vh - F'(x; h) = \mathcal{O}(\|h\|^2).$$

A semismooth function  $F : E^n \rightarrow E^n$  is *BD-regular* at  $x$  if all elements in  $\partial_B F(x)$  are nonsingular, and  $F$  is *strongly regular* at  $x$  if all elements in  $\partial F(x)$  are nonsingular.

# HOMOTOPY THEORY FOR NONSMOOTH FUNCTIONS

**Theorem.** Let  $F : E^n \rightarrow E^n$  be a Lipschitz continuous function and suppose there is a  $C^2$  map

$$\rho : E^m \times [0, 1) \times E^n \rightarrow E^n$$

such that

1.  $\nabla \rho(a, \lambda, x)$  has rank  $n$  on the set  $\rho^{-1}(\{0\})$ ,
2. the equation  $\rho_a(0, x) = 0$ , where  $\rho_a(\lambda, x) := \rho(a, \lambda, x)$ , has a unique solution  $x^a \in E^n$  for every fixed  $a \in E^m$ ,
3.  $\nabla_x \rho_a(0, x^a)$  has rank  $n$  for every  $a \in E^m$ ,
4.  $\rho$  is continuously extendible to the domain  $E^m \times [0, 1] \times E^n$ , and  $\rho_a(1, x) = F(x)$  for all  $x \in E^n$  and  $a \in E^m$ , and
5.  $\gamma_a$ , the connected component of  $\rho_a^{-1}(\{0\})$  containing  $(0, x^a)$ , is bounded for almost all  $a \in E^m$ .

Then for almost all  $a \in E^m$  there is a zero curve  $\gamma_a$  of  $\rho_a$ , along which  $\nabla \rho_a$  has rank  $n$ , emanating from  $(0, x^a)$  and reaching a zero  $\bar{x}$  of  $F$  at  $\lambda = 1$ . Further,  $\gamma_a$  does not intersect itself and is disjoint from any other zeros of  $\rho_a$ . Also, if  $\gamma_a$  reaches a point  $(1, \bar{x})$  and  $F$  is strongly regular at  $\bar{x}$ , then  $\gamma_a$  has finite arc length.

# SMOOTHING OPERATORS

**Definition.** Given a nonsmooth continuous function  $\phi : E^p \rightarrow E$ , a *smoother* for  $\phi$  is a continuous function  $\tilde{\phi} : E^p \times [0, \infty) \rightarrow E$  such that

1.  $\tilde{\phi}(x, 0) = \phi(x)$ , and
2.  $\tilde{\phi}$  is continuously differentiable on the set  $E^p \times (0, \infty)$ .

If  $\tilde{\phi}$  is  $C^2$  on  $E^p \times (0, \infty)$ , call  $\tilde{\phi}$  a  $C^2$ -*smoother*. For brevity, write  $\phi_\mu(x) := \tilde{\phi}(x, \mu)$ .

Example:  $\tilde{\phi}(a, b, \mu) := a + b - \sqrt{a^2 + b^2 + 2\mu}$  is a smoother for the NCP (nonlinear complementarity problem) function  $\phi(a, b) := a + b - \sqrt{a^2 + b^2}$ , where

$$\phi(a, b) = 0 \iff 0 \leq a \perp b \geq 0.$$

# APPLICATION TO COMPLEMENTARITY PROBLEMS

Nonlinear complementarity problem  $NCP(G)$ : find  $x \in E^n$  such that  $x \geq 0$ ,  $G(x) \geq 0$ ,  $x^t G(x) = 0$ .

Define  $F : E^n \rightarrow E^n$  by  $F_i(x) = \phi(x_i, G_i(x))$ ,  $F^\mu : E^n \rightarrow E^n$  by  $F_i^\mu(x) = \phi_\mu(x_i, G_i(x))$ . Then  $F^\mu$  is a smoother for  $F$ , and  $x$  solves  $NCP(G)$  if and only if

$$F(x) = 0.$$

Let  $\mu : [0, 1] \rightarrow [0, \infty)$  be a  $C^2$  decreasing function with  $\mu(1) = 0$ . Solve  $NCP(G)$  by using the homotopy map

$$\rho_a(\lambda, x) = \lambda F^{\mu(\lambda)}(x) + (1 - \lambda)(x - a).$$

## REFERENCES

- S. C. Billups and L. T. Watson, “A probability-one homotopy algorithm for nonsmooth equations and mixed complementarity problems”, *SIAM J. Optim.*, 12 (2002) 606–626.
- R. C. Melville, Lj. Trajković, S.-C. Fang, and L. T. Watson, “Artificial parameter homotopy methods for the DC operating point problem”, *IEEE Trans. Computer-Aided Design*, 12 (1993) 861–877.
- L. T. Watson, M. Sosonkina, R. C. Melville, A. P. Morgan, and H. F. Walker, “Algorithm 777: HOMPACT90: A suite of FORTRAN 90 codes for globally convergent homotopy algorithms”, *ACM Trans. Math. Software*, 23 (1997) 514–549.
- L. T. Watson, “Probability-one homotopies in computational science”, Technical Report TR 00-03, Dept. of Computer Sci., Virginia Polytechnic Institute and State Univ., Blacksburg, VA, 2000. <http://www.cs.vt.edu>
- S. M. Wise, A. J. Sommese, and L. T. Watson, “Algorithm 801: POLSYS\_PLP: A partitioned linear product homotopy code for solving polynomial systems of equations”, *ACM Trans. Math. Software*, 26 (2000) 176–200.